

Chiral bosons on Bargmann space associated with A_r Statistics

M. Daoud^{a1} and A. Hamama^b

^a *Max Planck Institute for Physics of Complex Systems,
Nöthnitzer Str. 38, D-01187 Dresden, Germany*

^b *High Energy Laboratory, Faculty of Sciences, University Mohamed V,
P.O. Box 1014 , Rabat , Morocco*

Abstract

We consider a large collection of particles obeying A_r statistics. The system behaves like a quantum droplet characterized by a constant Husimi distribution. We show that the excitations of this system live on the boundary of the droplet and they are described by an effective chiral boson action generalizing the Wess-Zumino-Witten theory in two dimension. Our analysis is based on the Fock-Bargmann analytical representations associated to A_r statistics. The quantization of the theory describing the dynamics on the edge is achieved. As by product, we prove that the edge excitations are given by a tensorial product of r abelian bosonic fields.

¹ *Faculté des Sciences, Département de Physique, Agadir, Morocco; email: m_daoud@hotmail.com*

1 Introduction

Fifty years ago, a first extension of Bose and Fermi statistics was achieved by Green [1]. This extension was the basic underlying mathematical background to investigate the implications of the generalizations of the familiar bosonic and fermionic statistics [2-6]. In the two last decades a renewal interest has been devoted to generalized quantum statistics due to their possible relevance in some issues like for instance fractional quantum Hall effect [7-8], anyon superconductivity [9] as well as black hole statistics [10]. Many variants of quantum statistics were proposed in the literature. One may quote the anyonic statistics [4] interpolating between fermionic and bosonic ones in two dimension space, quonic statistics [11] developed in the context of q -deformed algebras, k -fermionic statistics [12] defined as q -deformed version of ordinary bosons when the deformation parameter is such that $q^k = 1$ and Haldane fractional statistics [13] to explain the origin of the fractional quantization of Hall conductivity.

In the Green prescription [1], the generalized quantum statistics are characterized by certain triple relations which replace the commutation and anti-commutation rules for bosons and fermions. As by product, the fermions and bosons are promoted to para-fermions and para-bosons of order r respectively where the creation and annihilation operators satisfy

$$[[f_i^+, f_j^-], f_k^-] = -2\delta_{ik}f_j^-, \quad [[f_i^+, f_j^+], f_k^-] = -2\delta_{ik}f_j^+ + 2\delta_{jk}f_i^+, \quad [[f_i^-, f_j^-], f_k^-] = 0, \quad (1)$$

$$[\{b_i^+, b_j^-\}, b_k^-] = -2\delta_{ik}b_j^-, \quad [\{b_i^+, b_j^+\}, b_k^-] = -2\delta_{ik}b_j^+ - 2\delta_{jk}b_i^+, \quad [\{b_i^-, b_j^-\}, b_k^-] = 0, \quad (2)$$

with $i, j, k = 1, 2, \dots, r$. From an algebraic point of view, the Grassmann algebra in the fermionic case is replaced by the para-fermionic one (1) which is related to the orthogonal Lie algebra $so(2r+1) = B_r$ [14]. In the other side, the Weyl-Heisenberg algebra is extended to para-bosonic one (2) which is connected to the orthosymplectic superalgebra $osp(1/2r) = B(0, r)$ [15]. This indicates the deep link between Green statistics and the classical Lie and super Lie algebras. In this vain, very recently, on the basis of Palev works [6, 16], a classification of generalized quantum statistics were derived for the classical Lie algebras A_r , B_r , C_r and D_r [17].

This paper concerns the generalized A_r statistics. This generalization incorporates two kinds of statistics. The first one deals statistics satisfying a generalized exclusion Pauli principle and coincides with ones derived by Palev [6-16]. This class will be termed here fermionic A_r statistics. The second class is of bosonic kind. The particles can be accommodated in a given quantum state without any restriction. The generalization of A_r statistics was derived by one of the authors [18-20]. However, for completeness and in order to fix our notations we shall, in the next section, review the definition of the generalized A_r by means of the so-called Jacobson generators [21]. We use the trilinear relations defining the generalized quantum A_r statistics to construct the corresponding Fock space. We give the actions of the corresponding creation and

annihilation operators. The spectrum of the Hamiltonian, describing free particles obeying the generalized A_r statistics, is determined. The analytical Bargmann representations corresponding to Fock space of bosonic as well as fermionic A_r statistics are also presented. We give the differential realization of Jacobson operators. This analytical realization provides us with many advantages in discussing the semiclassical behavior of A_r quantum systems. Indeed, in the Bargmann space viewed as the phase space of A_r quantum systems, we study the excitations around a given droplet defined by a constant density (constant Husimi distribution). We show in this paper, that the excitations live on the boundary of the droplet and are described by an effective action generalizing Wess-Zumino-Witten one which describes chiral bosons in two dimensional space time [22]. To perform the semiclassical analysis of A_r quantum systems and the derivation of the effective theory of the edges excitations, some tools are needed like the star product in the Bargmann space. The necessary material to do this is presented. The strategy that we adopt is closer to one followed by Das et al [23] and Sakita [24] for non relativistic fermions localized around a ground many body state and forming a droplet. The droplet is specified by a diagonal density matrix ρ_0 . In the presence of an excitation potential, the fundamental state can be characterized by a unitary transformation of ρ_0 , namely $\rho_0 \longrightarrow U\rho_0U^\dagger$ where U is a collective variable describing the possible excitations around the droplet.

The paper is organized as follows. In section 2, a brief review of generalized A_r statistics, the associated Hamiltonian and the corresponding Fock space is given. The section 2 also deals with the Bargmann realization of the Jacobson generators (creation and annihilation operators of A_r quantum systems). This provides us with an useful way to analyze semi-classically the system under consideration. In this sense, we show in section 3 that a large collection of particles obeying A_r statistics behaves like a droplet in the Bargmann space. In this semi-classical description, the operators are replaced by functions and the commutators becomes Moyal brackets. In section 4, we show that the excitations of the system live on the boundary of the droplet and they are described by a chiral boson theory. The advantages of such formulation lies on the fact the dynamics of the obtained bosonic theory encodes the excitations of the A_r statistical system. It follows that our formulation provides a first convenient step to study large collective states of A_r statistics. The last section comprises concluding remarks.

2 The generalized A_r statistics

In this section, we introduce the definitions of the Jacobson operators and the generalized A_r statistics viewed as Lie triple system. We review the construction of the corresponding Fock space and we give the Hamiltonian describing a quantum system obeying generalized A_r statistics [18] (see also [19-20]).

2.1 Jacobson generators

First, let us introduce the notion of Lie triple system. Let V a vector space over a field F which is assumed to be either real or complex. The vector space V equipped vector with a trilinear mapping

$$[x, y, z] : V \otimes V \otimes V \longrightarrow V$$

is called Lie triple system if the following identities are satisfied:

$$[x, x, x] = 0,$$

$$[x, y, z] + [y, z, x] + [z, x, y] = 0,$$

$$[x, y, [u, v, w]] = [[x, y, u], v, w] + [u, [x, y, v], w] + [u, v, [x, y, w]].$$

According this definition, we will introduce the generalized A_r statistics as Lie triple system. In this respect, the algebra $\mathcal{G}$ defined by the generators a_i^+ and a_i^- ($i = 1, 2, \dots, r$) mutually commuting ($[a_i^-, a_j^-] = [a_i^+, a_j^+] = 0$) and satisfying the triple relation

$$[[a_i^+, a_j^-], x_k^+] = -s\delta_{jk}x_i^+ - s\delta_{ij}x_k^+ \quad (3)$$

$$[[a_i^+, a_j^-], a_k^-] = s\delta_{ik}x_j^- + s\delta_{ij}x_k^-, \quad (4)$$

where $s \in \{1, -1\}$, is closed under the ternary operation

$$[x, y, z] = [[x, y], z]$$

and define a Lie triple system. The elements $a_i^\pm$ are termed Jacobson generators and will be identified later with creation and annihilation operators of a quantum system obeying generalized A_r statistics. Note that for $s = -1$, the algebra $\mathcal{G}$ reduces to one defining the A_r statistics discussed in [6]. As we will see in what follows, the sign of the parameter s play an importance in the representation of the algebra $\mathcal{G}$ and consequently, one can obtain different microscopic and macroscopic statistical properties of the quantum system under consideration. Finally, recall that this statistics is intrinsically related to simple Lie algebras of class A like the para-fermion statistics which is related to class B of simple Lie algebras.

2.2 The Hamiltonian

The Jacobson generators $a_i^\pm$ can be identified with creation and annihilation operators of a quantum gas obeying the generalized A_r statistics. This requires a consistency with the Heisenberg equation

$$[H, a_i^\pm] = \pm e_i a_i^\pm \quad (5)$$

where H is the Hamiltonian of the system and the quantities e_i are the energies of the modes $i = 1, 2, \dots, r$. One can verify that if $|E\rangle$ is an eigenstate with energy E , $a_i^\pm|E\rangle$ are eigenvectors

of H with energies $E \pm e_i$. In this respect, the operators $a_i^\pm$ can be interpreted as ones creating or annihilating particles. To solve the consistency equation (5), we write the Hamiltonian H as

$$H = e_0 \mathbf{1} + h = e_0 \mathbf{1} + \sum_{i=1}^r e_i h_i \quad (6)$$

where e_0 is an arbitrary real constant and $\mathbf{1}$ is an operator commuting with all the elements of the algebra $\mathcal{G}$ (belonging to the centre of $\mathcal{G}$). The Hamiltonian H seems to be a simple sum of "free" (non-interacting) Hamiltonians h_i . However, it is important to note that in the quantum system under consideration, the statistical interactions occur and are encoded in the triple commutation relations (3) and (4). Using the structure relations of the algebra $\mathcal{G}$, the solution of the Heisenberg condition (5) is given by

$$h_i = \frac{s}{r+1} \left[(r+1)[a_i^-, a_i^+] - \sum_{j=1}^r [a_j^-, a_j^+] \right] + c \quad (7)$$

where the constant c will be defined later such that the ground state (vacuum) of the Hamiltonian H gives the energy e_0 .

2.3 Fock representations

An Hilbertian representation of the algebra $\mathcal{G}$ can be simply derived using the relations structures (3-4) defining A_r statistics. Here, we give the main results. For more details see the references [18,20]. Since, the algebra $\mathcal{G}$ is spanned by r pairs of Jacobson generators, it is natural to assume that the Fock space $\mathcal{F}$ is given by

$$\mathcal{F} = \oplus_{n=0}^{\infty} \mathcal{H}^n, \quad (8)$$

where

$$\mathcal{H}^n \equiv \{ |n_1, n_2, \dots, n_r\rangle, n_i \in \mathbf{N}, \sum_{i=1}^r n_i = n > 0 \}$$

and $\mathcal{H}^0 \equiv \mathbf{C}$. The action of $a_i^\pm$, on $\mathcal{F}$, are defined by

$$a_i^\pm |n_1, \dots, n_i, \dots, n_r\rangle = \sqrt{F_i(n_1, \dots, n_i \pm 1, \dots, n_r)} |n_1, \dots, n_i \pm 1, \dots, n_r\rangle \quad (9)$$

where the functions F_i are called the structure functions. Using the triple structure relations of A_r statistics, one obtains [18,20] the following expressions

$$F_i(n_1, \dots, n_i, \dots, n_r) = \frac{1}{2} n_i (2k - (1+s) + 2s(n_1 + n_2 + \dots + n_r)), \quad (10)$$

in terms of the quantum numbers $n_1, n_2, \dots, n_r$. In equation (10), the real parameter labelling the obtained representation satisfies the condition $2k - 1 > s$. The dimension of the irreducible representation space $\mathcal{F}$ is determined by the condition:

$$k - \frac{1+s}{2} + s(n_1 + n_2 + \dots + n_r) > 0. \quad (11)$$

It depends on the sign of the parameter s . It is clear that for $s = 1$, the Fock space $\mathcal{F}$ is infinite dimensional. However, for $s = -1$, there exists a finite number of basis states satisfying the condition $n_1 + n_2 + \cdots + n_r \leq k - 1$. The dimension is given, in this case, by $\frac{(k-1+r)!}{(k-1)!r!}$. This is exactly the dimension of the Fock representation of A_r statistics discussed in [6]. This condition-restriction is closely related to so-called generalized exclusion Pauli principle according to which no more than $k - 1$ particles can be accommodated in the same quantum state. In this sense, for $s = -1$, the generalized A_r quantum statistics give statistics of fermionic behavior. They will be termed here as fermionic A_r statistics and ones corresponding to $s = 1$ will be named bosonic A_r statistics. Having specified the Fock space associated with the generalized A_r statistics, one can obtain the spectrum of the Hamiltonian (6). For convenience, we set $c = \frac{2ks-s-1}{2r+2}$ in (7) and using the actions of creation and annihilation operators (9-10), one has

$$H|n_1, \cdots, n_i, \cdots, n_r\rangle = (e_0 + \sum_{i=1}^r e_i n_i)|n_1, \cdots, n_i, \cdots, n_r\rangle. \quad (12)$$

It is remarkable that, for $s = -1$, the spectrum of H is similar (with a slight modification) to energy eigenvalues of the A_r Calogero model (see for instance Eq.(1.2) in [25]). The latter describe the dynamical model containing $r + 1$ particles on a line with long rang interactions and provides a microscopic realization of fractional statistics [13,26]. It is important to stress that for $e_i = 0$ for all $i = 1, 2, \cdots, r$, the ground state energy e_0 is degenerate (the degeneracy coincides with the dimension of the Fock space; It is finite (resp. infinite) for fermionic (resp. bosonic) A_r statistics). It follows that $\hbar$ in equation (6) can be considered as a potential responsible of the degeneracy lifting and inducing fluctuations around the ground state energy e_0 . This remark constitutes the key ingredient, as it will be clarified later, to define A_r quantum droplets and to derive their excitations. Finally, we point out that for large k , we have

$$[a_i^-, a_j^+] \approx k\delta_{ij} \quad (13)$$

reflecting that the generalized A_r statistics (fermionic and bosonic ones) coincide with the Bose statistics and the Jacobson operators reduce to Bose ones (creation and annihilation operators of harmonic oscillators).

Besides the Fock representation discussed in this section, it is interesting to look for analytical realizations of the space representation associated with the Fock representations of the generalized A_r statistics. These realizations constitute an useful analytical tool in connection with variational and path integral methods to describe the quantum dynamics of the system described by the Hamiltonian H .

2.4 Bargmann realizations

First note that the Bargmann realization associated to A_r statistics was derived in [18] (see also [19-20]). Here, we recall some results needed for our task. This realization uses a suitably

defined Hilbert space of entire analytical functions. The Jacobson annihilation generators $a_i^\pm$ are realized as first order differential operators with respect to a complex variables z_i

$$a_i^- \longrightarrow \frac{\partial}{\partial z_i}. \quad (14)$$

The key point of such analytical realization lies on the fact that we represent the Fock states $|n_1, n_2, \dots, n_r\rangle$ as power of complex variables $z_1, z_2, \dots, z_r$:

$$|n_1, n_2, \dots, n_r\rangle \longrightarrow C_{n_1, \dots, n_r} z_1^{n_1} z_2^{n_2} \dots z_r^{n_r}. \quad (15)$$

Using the action of the annihilation operators on the Fock space $\mathcal{F}$ and the correspondences (14) and (15), the coefficients $C_{n_1, n_2, \dots, n_r}$ are obtained as

$$C_{n_1, n_2, \dots, n_r} = \left[\frac{(k-1+sn)!}{(k-1)!} \right]^{\frac{s}{2}} \frac{1}{\sqrt{n_1!} \sqrt{n_2!} \dots \sqrt{n_r!}} \quad (16)$$

where $n = n_1 + n_2 + \dots + n_r$ and $s = 1$ (resp. $s = -1$) for bosonic (resp. fermionic) A_r statistics. Using the equations (15-16), one can determine the differential action of the Jacobson creation operators. Indeed, from the actions of the generators a_i^+ on the Fock space and the triple relations (3) and (4), one obtains

$$a_i^+ \longrightarrow \frac{1}{2}(2k+s-1)z_i + sz_i \sum_{j=1}^r z_j \frac{d}{dz_j}. \quad (17)$$

The Jacobson generators act as first order linear differential operators. An arbitrary vector of the Fock space $\mathcal{F}$

$$|\phi\rangle = \sum_{n_1} \sum_{n_2} \dots \sum_{n_r} \phi_{n_1, n_2, \dots, n_r} |n_1, n_2, \dots, n_r\rangle,$$

is realized as

$$\phi(z_1, z_2, \dots, z_r) = \sum_{n_1} \sum_{n_2} \dots \sum_{n_r} \phi_{n_1, n_2, \dots, n_r} C_{n_1, n_2, \dots, n_r} z_1^{n_1} z_2^{n_2} \dots z_r^{n_r}. \quad (18)$$

The inner product of two functions ϕ and ϕ' is defined by

$$\langle \phi' | \phi \rangle = \int \int \dots \int d^2 z_1 d^2 z_2 \dots d^2 z_r \Sigma(k; z_1, z_2, \dots, z_r) \phi'^*(z_1, z_2, \dots, z_r) \phi(z_1, z_2, \dots, z_r) \quad (19)$$

The computation of the integration measure Σ , assumed to be isotropic, can be performed by choosing $|\phi\rangle = |n_1, n_2, \dots, n_r\rangle$ and $|\phi'\rangle = |n'_1, n'_2, \dots, n'_r\rangle$. A direct computation shows that the measure can be cast in the following compact form

$$\Sigma(\varrho_1, \varrho_2, \dots, \varrho_r) = \pi^{-r} \left[\frac{(k-1)!}{(k-sr+\frac{1}{2}(s-1))!} \right]^s [1 + s(\varrho_1^2 + \varrho_2^2 + \dots + \varrho_r^2)]^{sk-r-\frac{1}{2}(s+1)} \quad (20)$$

where $\varrho_i = |z_i|^2$. One can write the function $\phi(z_1, z_2, \dots, z_r)$ as the product of the state $|\phi\rangle$ with some ket $|\bar{z}_1, \bar{z}_2, \dots, \bar{z}_r\rangle$ labelled by the complex conjugate of the variables $z_1, z_2, \dots, z_r$

$$\phi(z_1, z_2, \dots, z_r) = \mathcal{N} \langle \bar{z}_1, \bar{z}_2, \dots, \bar{z}_r | \phi \rangle, \quad (21)$$

where $\mathcal{N}$ is a normalization constant to be adjusted later. Taking $|\phi\rangle = |n_1, n_2, \dots, n_r\rangle$, we have

$$\langle \bar{z}_1, \bar{z}_2, \dots, \bar{z}_r | n_1, n_2, \dots, n_r \rangle = \mathcal{N}^{-1} C_{n_1, n_2, \dots, n_r} z_1^{n_1} z_2^{n_2} \dots z_r^{n_r}. \quad (22)$$

This implies

$$|z_1, z_2, \dots, z_r\rangle = \mathcal{N}^{-1} \sum_{n_1} \sum_{n_2} \dots \sum_{n_r} \left[\frac{(k-1+sn)!}{(k-1)!} \right]^{\frac{s}{2}} \frac{z_1^{n_1}}{\sqrt{n_1!}} \frac{z_2^{n_2}}{\sqrt{n_2!}} \dots \frac{z_r^{n_r}}{\sqrt{n_r!}}. \quad (23)$$

It is important to notice that the expansion (23) converges for bosonic A_r statistics when $\bar{z}.z = |z_1|^2 + |z_2|^2 + \dots + |z_r|^2 < 1$. In other words, the complex variables $z_1, z_2, \dots, z_r$ should be in the complex domain defined by $\{(z_1, z_2, \dots, z_r) : |z_1|^2 + |z_2|^2 + \dots + |z_r|^2 < 1\}$. The normalization constant in (23) is given by

$$\mathcal{N} = (1 - s(|z_1|^2 + |z_2|^2 + \dots + |z_r|^2))^{-\frac{1}{4}(2ks-s+1)} \quad (24)$$

The states (23) are continuous in the labeling, constitute an over complete set in the respect to the measure given by (20) and then are coherent in the Klauder-Perelomov sense.

3 Semi classical analysis

One of the usefulness of the above Bargmann realizations is they provide us with a simply way to establish a correspondence between operators and classical functions on the phase space of the systems under consideration. So, in this section we shall investigate the semi classical properties of A_r statistical systems (bosonic as well as fermionic) in the Bargamnn space for k large.

3.1 The density matrix and Husimi distribution

It is commonly accepted that the exploration of the classical behavior of any quantum system hinges on whether one can describe the behavior of the wave-functions in terms of a density matrix. So, let $N = N_1 + N_2 + \dots + N_k$ the number of quantas of the system where N_i stands for the particle number in the mode i . The corresponding density operator is

$$\rho_0 = \sum_{n_1}^{N_1} \sum_{n_2}^{N_2} \dots \sum_{n_r}^{N_r} |n_1, n_2, \dots, n_r\rangle \langle n_1, n_2, \dots, n_r|. \quad (25)$$

In the Bargmann space, the mean value of the density matrix is defined by

$$\rho_0(\bar{z}, z) = \langle z | \rho_0 | z \rangle. \quad (26)$$

where z stands for the variables $(z_1, z_2, \dots, z_r)$ labelling the coherent states for A_r statistics systems. The mean value $\rho_0(\bar{z}, z)$ is the symbol associated with the density operator and it can be identified with the Husimi distribution for A_r quantum systems. As we are concerned with the situation when k is large, let us investigate the spacial shape of the mean value of the density operator. For A_r bosonic statistics, we have

$$\rho_0(\bar{z}, z) = (1 - \bar{z}.z)^k \sum_{n_1=0}^{N_1} \dots \sum_{n_r=0}^{N_r} \frac{(k-1+n_1+\dots+n_k)!}{(k-1)!n_1! \dots n_r!} |z_1|^{2n_1} \dots |z_k|^{2n_r} \quad (27)$$

It is easy to see that, for k large, the identity

$$(1 - \bar{z}.z)^{-k} = \sum_{n_1=0}^{\infty} \cdots \sum_{n_r=0}^{\infty} \frac{(k-1+n_1+\cdots+n_r)!}{(k-1)!n_1!\cdots n_r!} |z_1|^{2n_1} \cdots |z_r|^{2n_r}, \quad (28)$$

gives

$$(1 - \bar{z}.z)^k = \exp(-k\bar{z}.z). \quad (29)$$

Furthermore, using the relation

$$\sum_{n_1=0}^{N_1} \cdots \sum_{n_r=0}^{N_r} \frac{(k-1+n_1+\cdots+n_r)!}{(k-1)!n_1!\cdots n_r!} |z_1|^{2n_1} \cdots |z_r|^{2n_r} = \sum_{n=0}^N \frac{(k-1+n)!}{(k-1)!(n)!} (\bar{z}.z)^n \quad (30)$$

where $n = n_1 + \cdots + n_r$, one can see the term involving the sum in the expression of ρ_0 behaves like

$$\sum_{n=0}^N \frac{(k\bar{z}.z)^n}{n!}. \quad (31)$$

It follows that, for k large, the density can be approximated by

$$\rho_0(\bar{z}, z) \simeq \exp(-k\bar{z}.z) \sum_{n=0}^N \frac{(k\bar{z}.z)^n}{n!} \simeq \Theta(N - k\bar{z}.z) \quad (32)$$

for a large number N of particles. Clearly, $\rho_0(\bar{z}, z)$ is a step function for $k \rightarrow \infty$ and $N \rightarrow \infty$ ($\frac{N}{k}$ fixed). Similarly, the classical density for A_r fermionic statistics

$$\rho_0(\bar{z}, z) = (1 + \bar{z}.z)^{-\frac{k-1}{2}} \sum_{n_1=0}^{N_1} \cdots \sum_{n_r=0}^{N_r} \frac{(k-1)}{(k-1-n)!n_1!\cdots n_r!} |z_1|^{2n_1} \cdots |z_r|^{2n_r} \quad (33)$$

gives for large k and N

$$\rho_0(\bar{z}, z) \simeq \Theta(N - k\bar{z}.z). \quad (34)$$

It corresponds to a droplet configuration with boundary defined by $k\bar{z}.z = N$ and its radius is proportional to $\sqrt{N}$. The derivative of this density tends to a δ function. As we will see an interesting outcome of the semi-quantal dynamics happens when the parameter k tends to infinity. This indicates that k plays a crucial role in determining the classical limit of A_r quantum dynamics and in deriving the edge excitations for A_r fermionic as well as bosonic statistics.

3.2 The Star product and Moyal bracket

A second necessary ingredient to perform our semi classical analysis is the star product. In fact, as we will discuss next, for k large the mean value of the product of two operators leads to the Moyal star product. To show this, to every operator A acting on the Fock space $\mathcal{F}$, we associate the function

$$\mathcal{A}(\bar{z}, z) = \langle z|A|z \rangle. \quad (35)$$

An associative star product of two functions $\mathcal{A}(\bar{z}, z)$ and $\mathcal{B}(\bar{z}, z)$ is defined by

$$\mathcal{A}(\bar{z}, z) \star \mathcal{B}(\bar{z}, z) = \langle z|AB|z \rangle = \int d\mu(z', z') \langle z|A|z' \rangle \langle z'|B|z \rangle \quad (36)$$

where the measure $d\mu(\bar{z}, z) = d^2 z_1 d^2 z_2 \cdots d^2 z_r \Sigma$ is given by equation (20). To compute this star product, let us exploit the analytical properties of coherent states defined above. Indeed, Using the equations (23) and (24), one can see that the function defined by

$$\mathcal{A}(\bar{z}', z) = \frac{\langle z' | A | z \rangle}{\langle z' | z \rangle} \quad (37)$$

satisfy the following holomorphic and anti-holomorphic conditions

$$\frac{\partial}{\partial \bar{z}_i} \mathcal{A}(\bar{z}', z) = 0 \quad \frac{\partial}{\partial z'_i} \mathcal{A}(\bar{z}', z) = 0 \quad (38)$$

for $i = 1, 2, \dots, r$ and $z \neq z'$. Consequently, the action of the translation operator on the function $\mathcal{A}(\bar{z}', z)$ gives

$$\exp\left(z' \cdot \frac{\partial}{\partial z}\right) \mathcal{A}(\bar{z}', z) = \mathcal{A}(\bar{z}', z + z') \quad (39)$$

from which one can see that the function $\mathcal{A}(\bar{z}, z')$ is given by

$$\exp\left(-z \cdot \frac{\partial}{\partial z'}\right) \exp\left(z' \cdot \frac{\partial}{\partial z}\right) \mathcal{A}(\bar{z}, z) = \exp\left((z' - z) \cdot \frac{\partial}{\partial z}\right) \mathcal{A}(\bar{z}, z) = \mathcal{A}(\bar{z}, z') \quad (40)$$

in term of the function $\mathcal{A}(\bar{z}, z)$. Similarly, one obtains

$$\exp\left(-\bar{z} \cdot \frac{\partial}{\partial \bar{z}'}\right) \exp\left(\bar{z}' \cdot \frac{\partial}{\partial \bar{z}}\right) \mathcal{A}(\bar{z}, z) = \mathcal{A}(\bar{z}', z) \quad (41)$$

Equivalently, the equations (40) and (41) can also be cast in the following forms

$$\exp\left((z' - z) \cdot \frac{\partial}{\partial z}\right) \mathcal{A}(\bar{z}, z) = \mathcal{A}(\bar{z}, z') \quad (42)$$

and

$$\exp\left((\bar{z}' - \bar{z}) \cdot \frac{\partial}{\partial \bar{z}}\right) \mathcal{A}(\bar{z}, z) = \mathcal{A}(\bar{z}', z) \quad (43)$$

respectively. Combining the equations (36-37) and (42-43), the star product rewrites as

$$\mathcal{A}(\bar{z}, z) \star \mathcal{B}(\bar{z}, z) = \int d\mu(\bar{z}', z') \exp\left((z' - z) \cdot \frac{\partial}{\partial z}\right) \mathcal{A}(\bar{z}, z) |\langle z | z' \rangle|^2 \exp\left((\bar{z}' - \bar{z}) \cdot \frac{\partial}{\partial \bar{z}}\right) \mathcal{B}(\bar{z}, z) \quad (44)$$

where the overlapping of coherent states is given by

$$\langle z | z' \rangle = \left[(1 - s \bar{z}' \cdot z') (1 - s \bar{z} \cdot z) (1 - s \bar{z}' \cdot z)^{-2} \right]^{\frac{2ks+1-s}{4}} \quad (45)$$

with $s = +1, -1$ corresponding to bosonic and fermionic statistics respectively. Clearly, the modulus of the kernel (45) possesses the properties $|\langle z | z' \rangle| = 1$ if and only if $z = z'$, $|\langle z | z' \rangle| < 1$ and $|\langle z | z' \rangle| \rightarrow 0$ for k large. The latter properties are helpful to get the star product between two functions on the Bargmann space. In this respect, we introduce a function $s(z', z)$ of the coordinates of two points on the Bargmann space

$$s^2(z', z) = -\ln |\langle z | z' \rangle|^2 = \frac{1}{2} (2ks - s + 1) \ln \frac{(1 - s \bar{z}' \cdot z) (1 - s \bar{z} \cdot z')}{(1 - s \bar{z} \cdot z) (1 - s \bar{z}' \cdot z')}. \quad (46)$$

It verifies the properties : $s(z', z) = s(z, z')$ and $s(z', z) = 0$ if and only if $z' = z$. This function can be interpreted as the distance between two points on the Bargmann space. It turns out that the overlapping (45) generates the metric. In fact the line element ds^2 , defined as the quadratic part of the decomposition of $s^2(z, z + dz)$ (distance between two infinitesimal points), is given by

$$ds^2 = g_{i\bar{j}} dz_i d\bar{z}_j \quad (47)$$

where summation over repeated indices is understood and the components of the metric $g_{i\bar{j}}$ are defined

$$g_{i\bar{j}} = (k + \frac{s}{2} - \frac{1}{2}) \left[\frac{\delta_{ij}}{1 - s\bar{z}.z} + s \frac{\bar{z}_i z_j}{(1 - s\bar{z}.z)^2} \right]. \quad (48)$$

We now come to the evaluation of the star product for k large. Since $s^2(z', z)$ tends to infinity with $k \rightarrow \infty$ if $z \neq z'$ and equals zero if $z = z'$, one can conclude that, in that limit, the domain $z \simeq z'$ gives only a contribution to the integral (44). Decomposing the integrand near the point $z \simeq z'$ and going to integration over $\eta = z' - z$, one gets

$$\mathcal{A}(\bar{z}, z) \star \mathcal{B}(\bar{z}, z) = \int \frac{d\eta. d\bar{\eta}}{\pi^r} \exp\left(\eta. \frac{\partial}{\partial z}\right) \mathcal{A}(\bar{z}, z) \exp\left(-g_{i\bar{j}} \eta_i \bar{\eta}_j\right) \exp\left(\bar{\eta}. \frac{\partial}{\partial \bar{z}}\right) \mathcal{B}(\bar{z}, z). \quad (49)$$

It follows that the star product between two functions on the Bargmann space associated with A_r statistics is given by

$$\mathcal{A}(\bar{z}, z) \star \mathcal{B}(\bar{z}, z) = \mathcal{A}(\bar{z}, z) \mathcal{B}(\bar{z}, z) - g^{i\bar{j}} \frac{\partial \mathcal{A}}{\partial z_i}(\bar{z}, z) \frac{\partial \mathcal{B}}{\partial \bar{z}_j}(\bar{z}, z) + O\left(\frac{1}{k^2}\right) \quad (50)$$

where the matrix

$$g^{i\bar{j}} = 2 \frac{1 - s\bar{z}.z}{2k + s - 1} (\delta_{ij} - s z_i \bar{z}_j) \quad (51)$$

is the inverse of the metric $g_{i\bar{j}}$ and it is proportional to $1/k$. Then, the symbol or function associated with the commutator of two operators A and B is given by

$$\langle z|[A, B]|z \rangle = \{\mathcal{A}(\bar{z}, z), \mathcal{B}(\bar{z}, z)\}_\star = -g^{i\bar{j}} \left(\frac{\partial \mathcal{A}}{\partial z_i}(\bar{z}, z) \frac{\partial \mathcal{B}}{\partial \bar{z}_j}(\bar{z}, z) - \frac{\partial \mathcal{B}}{\partial z_i}(\bar{z}, z) \frac{\partial \mathcal{A}}{\partial \bar{z}_j}(\bar{z}, z) \right) \quad (52)$$

where

$$\{\mathcal{A}(\bar{z}, z), \mathcal{B}(\bar{z}, z)\}_\star = \mathcal{A}(\bar{z}, z) \star \mathcal{B}(\bar{z}, z) - \mathcal{B}(\bar{z}, z) \star \mathcal{A}(\bar{z}, z). \quad (53)$$

is the so-called the Moyal bracket.

3.3 The excitation potential

The quantum droplet under consideration is specified by the density matrix ρ_0 (25). The excitations of this configuration can be described by a unitary time evolution operator U which gives information concerning the dynamics of the excitations around ρ_0 . The excited states will be characterized by a density operator $\rho = U \rho_0 U^\dagger$. In this respect, the Hamiltonian h in equation (6) may be viewed as the excitation potential of the quantum droplet. Indeed, as we mentioned in the previous section, in absence of h (all e_i vanishing) the states $|n_1, n_2, \dots, n_r\rangle$

are eigenstates of H with the same eigenvalue e_0 . The degeneracy of the energy e_0 coincides with the dimension of the Fock space $\mathcal{F}$. It is finite for A_r bosonic systems and takes a finite value for A_r fermionic statistics. The Hamiltonian h is exactly the excitation potential that induces a degeneracy lifting. Using the expressions (12) and (23), the mean value of the excitation potential h is

$$\langle z|h|z\rangle = \mathcal{H}(\bar{z}, z) = (k + \frac{s}{2} - \frac{1}{2}) \sum_{i=1}^r e_i \frac{z_i \bar{z}_i}{1 - s \bar{z} \cdot z}. \quad (54)$$

The function $\mathcal{H}(\bar{z}, z)$ is the symbol associated to h . As we will next concerned by the edge excitations living on the boundary of the A_r quantum droplet, it is simply verified from equations (23) and (24) that the Hamiltonian symbol $\mathcal{H}$ takes for k large the simple form

$$\mathcal{H}(\bar{z}, z) = k \sum_{i=1}^r e_i z_i \bar{z}_i, \quad (55)$$

which is just the classical harmonic oscillator potential.

4 Edge excitations and chiral bosons action

4.1 Effective action

In this section, we derive the effective action for excitations living on the edge of a A_r quantum droplet. The derivation is based on semi classical analysis given in the previous section. As mentioned above, the dynamical information, related to degrees of freedom of the edge states, is contained in the unitary operator U . The corresponding action is

$$S = \int dt \text{Tr}(\rho_0 U^\dagger (i\partial_t - H) U). \quad (56)$$

It is compatible with the Liouville evolution equation for the density matrix

$$i \frac{\partial \rho}{\partial t} = [H, \rho].$$

To write down an effective action describing the edge excitations, we evaluate the quantities occurring in (56) as classical functions on the basis of the semi classical analysis performed above. We start by computing the term $i \int dt \text{Tr}(\rho_0 U^\dagger \partial_t U)$. For this, we set $U = e^{+i\Phi}$ ($\Phi^\dagger = \Phi$). A direct computation gives

$$dU = \sum_{n=1}^{\infty} \frac{(i)^n}{n!} \sum_{p=0}^{n-1} \Phi^p d\Phi \Phi^{n-1-p}, \quad (57)$$

from which one obtains

$$U^\dagger dU = i \int_0^1 d\alpha e^{-i\alpha\Phi} d\Phi e^{+i\alpha\Phi}. \quad (58)$$

Thus, we have

$$e^{-i\Phi} \partial_t e^{+i\Phi} = i \int_0^1 d\alpha e^{-i\alpha\Phi} \partial_t \Phi e^{+i\alpha\Phi}. \quad (59)$$

Using Baker-Campbell-Hausdorff formula, one can show

$$i \int dt Tr(\rho_0 U^\dagger \partial_t U) = \int dt \sum_{n=0}^{\infty} \frac{-(i)^n}{(n+1)!} Tr(\underbrace{[\Phi, \dots [\Phi, \rho_0] \dots]_n}_{n} \partial_t \Phi) \quad (60)$$

Due to the coherent states completeness, the trace of any operator A is

$$Tr A = \int d\mu(\bar{z}, z) \langle z | A | z \rangle.$$

It follows that the equation (60) rewrites as

$$i \int dt Tr(\rho_0 U^\dagger \partial_t U) = \int d\mu dt \sum_{n=0}^{\infty} \frac{-(i)^n}{(n+1)!} \underbrace{\{\Phi, \dots \{\Phi, \rho_0\}_\star \dots\}_\star}_n \star \partial_t \Phi \quad (61)$$

where the star product and the Moyal bracket are respectively defined by (50) and (52). It is important to stress that ρ_0 and Φ in equation (61) are now classical functions. It is easy to see that the equation (61) gives

$$i \int dt Tr(\rho_0 U^\dagger \partial_t U) \simeq -\frac{i}{2} \int d\mu dt \{\Phi, \rho_0\}_\star \partial_t \Phi \quad (62)$$

where we have dropped terms in $\frac{1}{k^2}$ as well as as the total time derivative. Using the expression (52), the Moyal bracket in (62) writes

$$\{\Phi, \rho_0\}_\star = \frac{2i}{2k + s - 1} (\mathcal{L}\Phi) \frac{\partial \rho_0}{\partial(\bar{z}.z)} \quad (63)$$

where the first order differential operator is

$$\mathcal{L} = i(1 - s\bar{z}.z)^2 (z \cdot \frac{\partial}{\partial z} - \bar{z} \cdot \frac{\partial}{\partial \bar{z}}). \quad (64)$$

For k large the density function (see equations (32) and (34)) is a step function. Its derivative is a δ function with support on the boundary $\partial\mathcal{D}$ of the droplet $\mathcal{D}$ defined by $k\bar{z}.z = N$. It follows

$$i \int dt Tr(\rho_0 U^\dagger \partial_t U) \approx -\frac{1}{2} \int d\mu dt \delta(N - k\bar{z}.z) (\mathcal{L}\Phi) (\partial_t \Phi) = -\frac{1}{2} \int_{\partial\mathcal{D} \times \mathbf{R}^+} dt (\mathcal{L}\Phi) (\partial_t \Phi) \quad (65)$$

The second step in the derivation of edge states action consists in the simplification the second term in (56) involving H . By a straightforward calculation, we obtain

$$Tr(\rho_0 U^\dagger h U) = Tr(\rho_0 h) + i Tr([\rho_0, h] \Phi) + \frac{1}{2} Tr([\rho_0, \Phi] [h, \Phi]) \quad (66)$$

The first term in r.h.s of (66) is Φ -independent. We drop it since it no contains any information about the dynamics of the edge excitations. The second term in r.h.s of (66) rewrites as

$$i Tr([\rho_0, h] \Phi) \approx i \int d\mu \{\rho_0, \mathcal{H}\}_\star \Phi \quad (67)$$

in term of the Moyal bracket where $\mathcal{H}$ is given by (55). By a direct computation, one can see that

$$i Tr([\rho_0, h] \Phi) \longrightarrow 0 \quad (68)$$

The last term in r.h.s of (66) gives

$$\frac{1}{2}Tr([\rho_0, \Phi][h, \Phi]) \approx \frac{i}{2k} \int d\mu dt (\mathcal{L}\Phi) \frac{\partial \rho_0}{\partial(\bar{z}.z)} \{\mathcal{H}, \Phi\}_\star \quad (69)$$

where the Moyal bracket is given by

$$\{\mathcal{H}, \Phi\}_\star = \frac{2k(1 - s\bar{z}.z)}{2k + s - 1} \left[s(z.\frac{\partial \Phi}{\partial z} - \bar{z}.\frac{\partial \Phi}{\partial \bar{z}}) \sum_{i=1}^r e_i \bar{z}_i z_i - \sum_{i=1}^r e_i (z_i \frac{\partial \Phi}{\partial z_i} - \bar{z}_i \frac{\partial \Phi}{\partial \bar{z}_i}) \right] \quad (70)$$

Since the derivative of the density ρ_0 gives a delta function with support on the boundary of the quantum droplet, the equation (69) is simplified

$$\{\mathcal{H}, \Phi\}_\star \approx i \sum_{i=1}^r e_i \mathcal{L}_i \Phi \quad (71)$$

where $\mathcal{L}_i$ is the angular momentum with respect to the variable z_i (see the definition (64)).

Finally, we obtain

$$\int dt Tr(\rho_0 U^\dagger H U) = \frac{1}{2} \int d\mu dt \delta(N - k\bar{z}.z) (\mathcal{L}\Phi) \sum_{i=1}^r e_i (\mathcal{L}_i \Phi) + O(\frac{1}{k^2}) \quad (72)$$

Note that we have eliminated the term containing the ground states energy e_0 which does not contribute to the edge dynamics. Combining (65) and (72), we get

$$S \approx -\frac{1}{2} \int_{\partial \mathcal{D} \times \mathbf{R}^+} d\mu dt \delta(N - k\bar{z}.z) (\mathcal{L}\Phi) \left((\partial_t \Phi) + \sum_{i=1}^r e_i (\mathcal{L}_i \Phi) \right). \quad (73)$$

This action involves only the time derivative of Φ and the tangential derivatives ($\mathcal{L}_i \Phi$). It is a generalization of a chiral abelian Wess-Zumino-Witten (WZW) theory [22]. It is interesting to note that for $r = 1$, we recover the WZW action describing a bosonized theory of a system of large fermions in two dimension [25]. Solving the equations of motion arising from the action (73) gives the nature of edge states. This will be done in the next subsection.

4.2 Edge fields

The action (73) is minimized by the fields Φ that satisfy the equation of motion

$$\mathcal{L}(\partial_t \Phi + \sum_{i=1}^r e_i \mathcal{L}_i \Phi) = 0. \quad (74)$$

Since the theory is defined on the boundary of the droplet fixed by the condition $\bar{z}.z = \frac{N}{k}$, we introduce the angular variables θ_i ($z_i = \sqrt{\frac{N}{k}} e^{i\theta_i}$). This is the most simple parametrization that one can consider. The operators $\mathcal{L}_i$ reduce to partial derivatives ∂_i with respect to θ_i . The general solution of the equation of motion can be written as

$$\Phi(\theta_1, \theta_2, \dots, \theta_r, t) = \Phi(\theta_1 - e_1 t, \theta_2 - e_2 t, \dots, \theta_r - e_r t) + \Lambda(t). \quad (75)$$

In the last equation, $\Lambda(t)$ represents the gauge degree of freedom corresponding to the invariance of the action (73) under the transformation

$$\Phi \longrightarrow \Phi + \Lambda(t).$$

This can be discarded by imposing the gauge condition

$$(\partial_t + \sum_{i=1}^r e_i \partial_i) \Phi = 0. \quad (76)$$

Next, we assume that the field Φ can be expressed in a factorized form

$$\Phi = \Phi_1 \Phi_2 \cdots \Phi_r \quad (77)$$

in terms of r components $\Phi_i = \Phi_i(\theta_i, t)$ ($i = 1, 2, \dots, r$) satisfying the equations

$$(\partial_t + e_i \partial_i) \Phi_i = 0. \quad (78)$$

It is remarkable that the equations (77) and (78) are compatible with the gauge fixing condition (76). To obtain the solution of (78), we assume that the field Φ_i satisfies the following periodicity condition

$$\Phi_i(2\pi, t) - \Phi_i(0, t) = -2\pi \alpha_0^i \quad (79)$$

where α_0^i is a time independent constant. It is easy to see that the general solution of (78) is then given by

$$\Phi_i(\theta_i, t) = \bar{\alpha}_0^i - \alpha_0^i(\theta_i - e_i t) + i \sum_{n \neq 0} \frac{\alpha_n^i}{n} e^{in(\theta_i - e_i t)} \quad (80)$$

where the constant $\bar{\alpha}_0^i$ can be viewed as the canonical momentum associated to α_0^i . Note also that the complex coefficients in (80) satisfy $(\alpha_n^i)^* = \alpha_{-n}^i$ required by the reality condition of the field Φ_i . The canonical momentum corresponding to the field Φ_i is

$$\Pi_i(\theta_i, t) = \alpha_0^i + \sum_{n \neq 0} \alpha_n^i e^{in(\theta_i - e_i t)}. \quad (81)$$

The quantization of the theory of edge excitations described by the action (73) can be performed by imposing the equal time commutation rules

$$[\Pi_i(\theta_i, t), \Phi_j(\theta_j, t)] = i \delta_{ij} \delta(\theta_i - \theta_j). \quad (82)$$

This implies that $\bar{\alpha}_0^i$, α_0^i and α_n^i become operators satisfying the relations

$$[\alpha_n^i, \alpha_m^j] = \delta_{ij} \delta_{m+n,0} \quad [\alpha_0^i, \bar{\alpha}_0^j] = i \delta_{ij}. \quad (83)$$

The other commutators vanish. This reflects that each field Φ_i is a superposition of oscillating modes on the boundary of the A_r droplet. For a fixed i , the Hilbert space $\mathcal{H}_i$ is a tensorial product of harmonic oscillator Fock spaces. The whole Hilbert space is then given by

$$\mathcal{H} = \mathcal{H}_1 \otimes \mathcal{H}_2 \cdots \otimes \mathcal{H}_r. \quad (84)$$

5 Concluding remarks

To conclude, let summarize the main points discussed in this paper. We provided a general approach of what we agreed to call " Chiral boson theory on Bargmann space associated with A_r quantum statistics". We started discussing the essential structures of these new quantum statistics. We also gave the analytical Bargmann realizations. In the Bargmann space, the problem of computing commutators has been rephrased in terms of (more easy) Moyal brackets of functions associated to the algebra generated by creation and annihilation (Jacobson) operators. Moyal bracket captures the essence of the full quantum characteristics of A_r statistics. This potentially provides us with an important tool to obtain semi-classically the effective action describing a large collection of N particles with A_r statistics. More precisely, we have shown that for large N and large k (k the parameter indexing the Fock representations), the system behaves like a droplet in the Bargmann space (phase space). We derived the effective action (equation (73)) describing the excitations living on the droplet's boundary. It is remarkable that the obtained action is similar to Wess-Zumino-action [22]. As by product, we have shown that the boundary excitations are essentially a tensorial product of r bosonic fields (cf. equation (77)). Each bosonic field is given in terms of an infinite oscillating modes (harmonic oscillators) (see equation (80)).

The results of the present paper can be used in connection with the so-called generalized spin systems [27]. In fact, the usual $SU(2)$ spin system are extended to spin models based on an arbitrary Lie group. In particular, for the classical Lie algebras of class A_r , the $SU(2)$ spin generators are replaced by the A_r generators which coincide with the creation and annihilation (Jacobson) operators in the terminology of this paper. It follows that the generalized spin models, discussed in [27], can be viewed as example of systems with generalized A_r statistics. In this sense, we believe that the approach developed here can be adapted to the generalized spin systems. We hope to report on this issue in a forthcoming work.

Acknowledgments: One of us (M. D.) would like to thank the Max Planck Institute for Physics of Complex Systems for the kind hospitality extended to him during December 2007.

References

- [1] H.S. Green, Phys. Rev. **90** (1953) 270.
- [2] O.W. Greenberg and A.M.L. Messiah, Phys. Rev. B **138** (1965) 1155;
J. Math. Phys. **6** (1965) 500.
- [3] Y. Ohnuki and S. Kamefuchi, *Quantum field theory and parastatistics*
(Springer, Berlin, 1982).
- [4] J.M. Leinaas and J. Myrheim, Nuovo Cimento B **37** (1977) 1.
- [5] F. Wilczek, Phys. Rev. Lett. **49** (1982) 957.

- [6] T.D. Palev, *Lie algebraic aspects of quantum statistics. Unitary quantization (A-quantization)*, Preprint JINR E17-10550 (1977) and hep-th/9705032.
- [7] R.B. Laughlin, Phys. Rev. Lett. **50** (1983) 1395.
- [8] B.I. Halperin, Phys. Rev. Lett. **52** (1984) 1583.
- [9] R.B. Laughlin, Phys. Rev. Lett. **60** (1988) 2677.
- [10] A. Strominger, Phys. Rev. Lett. **71** (1993) 3397.
- [11] O.W. Greenberg, Phys. Rev. Lett. **64** (1990) 705; Phys. Rev. D **43**(1991) 4111.
- [12] M. Daoud and M. Kibler, Phys. Lett. A **321** (2004) 147; Phys. Part. Nuclei (Suppl. 1) **33** (2002) S43.
- [13] F.D.M. Haldane, Phys. Rev. Lett. **67** (1991) 937.
- [14] S. Kamefuchi and Y. Takahashi, Nucl. Phys. **36** (1962) 177; C. Ryan and E.C.G. Sudarshan, Nucl. Phys. **47** (1963) 207.
- [15] V.G Kac, Adv. Math. **26** (1977) 8.
- [16] T.D. Palev, Czech. J. Phys. **B 29** (1979) 91; Rep. Math. Phys. **18** (1980) 177; Rep. Math. Phys. **18** (1980) 129; J. Math. Phys **21** (1980) 1293.
- [17] N.I. Stoilova and J. Van der Jeugt, J. Math. Phys. **46** (2005) 033501.
- [18] M. Daoud, J. Phys. A: Math. Gen. **39** (2006) 889.
- [19] M. Daoud, Inter. Jour. of Mod. Phys. A **21** (2006) 4543.
- [20] M. Daoud, Mod. Phys. Lett. A **21** (2006) 1691.
- [21] N. Jacobson, Amer. J. Math. **71** (1949) 149.
- [22] E. Witten, Comm. Math. Phys. **92** (1984) 455.
- [23] S. R. Das, A. Dhar, G. Mandal and S. R. Wadia, Int. J. Mod. Phys. **A 7** (1992) 5165; Mod. Phys. Lett. **A 7** (1992) 71.
- [24] B. Sakita, Phys. Lett. B. **387** (1996) 118.
- [25] B. Basu-Mallick, Int. J. Mod. Phys. B **16** (2002) 1875.
- [26] A. P. Polychronakos, in *Les Houches Lectures* (1998), hep-th/9902157.
- [27] S. Randjbar-Daemi, Abdus Salam and J. Strathdee, Phys. Rev. B **48** (1993) 3190.